SUBJECT INDEX - Volumes 1 and 2

acute hepatitis viral infection, pathogenesis and serodiagnosis of, 2, 31

adeno-associated virus, growth in embryonated eggs, 2, 287

adenovirus

detection in stool specimens by EIA, 1, 331 neutralization of cytotoxin, in Vero cells, 2, 321

rapid identification of, 2, 367

affinity chromatography

use in preparation of erythrocyte receptors, 1, 285

use to purify bovine diarrhea virus, 2, 293 use to purify reverse transcriptase from avian retroviruses, 1, 157

African green monkey kidney cell line, for production of adeno-associated virus, 2, 287

agarose gel electrophoresis, separation of M-protein from influenza virus by, 2, 211

aldehyde fixation, of erythrocytes, for coupling to HBsAg, 2, 269

alkali-liberated nuclear polyhedrosis virus, gipsy moth, field-flow fractionation, 1, 241

anti-complement immunoperoxidase, compared to ACIP and ACIF methods for detection of Epstein-Barr virus, 1, 11

antigenic priming, with influenza virus, previous, detection of, 1, 355

anti-HBc clinical relevance of, 2, 107

detection and significance in the Blood Bank, 2, 119

detection by RIA and counterimmunoelectrophoresis, 2, 181

in patients with hepatitis B, 2, 97

anti-HBe

EIA for, 2, 85

in patients with hepatitis B, 2, 97

anti-HBs

clinical relevance of, 2, 107 confirmation of specificity in immuno-

radiometric assay, 1, 113 EIA for rapid detection of, 2, 63

hybridoma cells secreting, 1, 257

markers for in immune electron microscopy, 2, 169

method for coupling erythocytes for use in passive haemagglutination, 2, 269

screeining of blood donors/recipients for by EIA, 2, 71

sensitive radiometric assay for detection of, 1, 311

anti-IgG immunoperoxidase, compared to ACIP and ACIF methods for the detection of Epstein-Barr virus, 1, 11

anti-IgM conjugate, to detect rheumatoid factor, 1, 1

antigenic priming, previous, detected by plaque inhibition test, 1, 355

antigens

associated with long-incubation non-A, non-B hepatitis virus, identification and detection of, 2, 127

location and quantitation of on the surface of virus-infected cells, 1, 61

of rabies virus origins, identification of in brain impression smears, 1, 33

antiviral antibodies, hen egg yolk as source of to plant viruses in ELISA, 1, 179

arboviruses

effect of some disinfectants on, 1, 27

simple serological diagnosis of, 1, 299 avian adenovirus, used as helper for preparing adeno-associated virus, 2, 287

avian myeloblastosis virus, purified using affinity chromatography, 1, 157

avian sarcoma virus, size determination of by antibody-mediated polysome precipitation, 1, 99

B.95-9 cells, continuous culturing of in the presence of phosphonoacetic acid, 1, 349 bacterial adherence, specific, for location and

quantitation of antigen on the surface of virus-infected cells, 1,61

bacteriophage

lysates, concentration of by filter chromatography, 1, 79

rapid concentration of from large volumes of freshwater, 1, 87

 ϕ X174, electron microscopic studies of, 2, 159

Blood Bank, detection and significance of anti-HBc and HBcAg in the, 2, 119

blood products, administration of, followed by development of non-A, non-B hepatitis, 2, 17

Bombyx mari (silkworm), use of electron microscopic studies of bacteriophage φX174, intact and 'eclipsing' particles, 2, 159

bovine diarrhea virus, affinity chromatography of, 2, 293

canine kidney cells, transformation of by simian virus 40, 2, 331

cell-mediated cytotoxic responses, ³H-uridine used to measure, 2, 237

chicken embryo lethal orphan virus (CELO), used as helper for preparing adeno-associated virus, 2, 287

chromosome—EBNA interaction, studied by use of new technique for detection of Epstein—Barr nuclear antigen or anti-EBNA antibodies, *I*, 11

chronic liver disease

persistence of anti-HBc IgM in, 2, 97

progression to by cases of acute viral hepatitis and development of in renal and hepatic transplant recipients, 2, 17

citrus exocortis virus, use in hybridisation studies, 1, 229

citrus tristeza virus (CTV), antibodies against raised in hen egg yolk, 1, 179

coliphages MS-2 and T2, concentration of by filter chromatography, 1, 79

complementary DNA, in immunochemical detection of ds RNA in plants by hybridisation of viral ds RNA to, 2, 277

complement fixation

single radial, 2, 261

test for assay of specimens for HAV or anti-HAV, 2, 31 congenital cytomegalovirus infection in newborns, routine screening for, 2, 315

counterimmunoelectrophoresis

*comparison of with RIA for the detection of anti-HBc, 2, 181

use for identification of protein fractions, 2, 211

Coxsackie virus rapidly identified by indirect immune electron microscopy technique, 2, 367

Crotalaria juncea, lectin isolated from, used in affinity chromatography, 2, 293

C-type viral glycoproteins, ELISA for detection and quantification of, 2, 357

CVS, rabies strains, identified by use of monoclonal antibodies, 1, 33

cytomegalovirus

antibodies, detection and differentiation of by RIA, 1, 133

antigen and antibody, interchangeable ELISA for 2, 301, 315

congenital infection, routine screeining of newborns with, 2, 315

cytotoxin, adenovirus, neutralisation of, 2, 321

Dane particles, characterisation of, 1, 47 diarrhea (bovine viral), 2, 293 disinfectant testing (virucide), 1, 27 DNA

complementary, in immunochemical detection of ds RNA in plants, 2, 277

orthopox, preparation of from viral cores, 2, 175

DNA polymerase

hepatitis B virus, marker of virus replication, 2, 7

hepatitis B virus-specific, technique for the detection of, 2, 349

dye uptake and dilution technique, as a method for measuring growth inhibition by interferon, 1, 197

'eclipsing' particles, electron microscopic studies of, 2, 159

electron microscopy

congenital infection, routine screening of newborns with, 2, 315

in the characterisation of rubella virus revealing haemagglutinin rosettes, 2, 195 rotavirus, examined by for markers, 2, 169 studies of bacteriophage ϕ X174 intact and 'eclipsing' particles, 2, 159

electrophoresis, in the detection of replicating, super-coiled simian virus 40 DNA, I, 235

embryonated eggs, growth of adeno-associated virus in, 2, 287

embryonic kidney cells (human), somatic cell hybrids constructed with, for study of viral gene expression, *I*, 139

encephalocarditis virus, preparation of erythrocyte receptors for by affinity chromatography, 1, 285

enzyme-immunoassay (EIA)

compared with RIA for detection of rotaviruses and adenoviruses, 1, 331

for HBeAg of enzyme-labelled F(ab')₂ and IgG conjugates, 2, 85

for rapid detection of HBsAg and anti-HBs, 2,63

Hepanostika, for HBsAg, improved, 2, 57 use in testing for HBsAg and surface antibody, 2, 71

enzyme-labelled F(ab')₂ compared with IgG conjugates in an EIA for HBeAg, 2, 85 enzyme-linked immunosorbent assay (ELISA) for detection and quantification of C-type viral glycoproteins, 2, 357

for establishing diagnosis of hepatitis A, 2, 47 hen egg yolk as source of antiviral antibodies to plant viruses in, 1, 179

interchangeable, for cytomegalovirus antigen and antibody, 2, 301, 315

or rubella IgM and IgG antibodies, interference of IgM rheumatoid factor in, 1, 1 epidemiology, of hepatitis A in Amsterdam, 2, 47

Epstein-Barr virus

presence of tested for in lymphoblastoid B.95-8 cells, 1, 349

technique for the detection of, 1, 11

erythrocyte receptors, for viruses, preparation of by affinity chromatography, 1, 285 erythrocytes

aldehyde-fixed, coupled to HBsAg for use in passive haemagglutination, 2, 269 formalinised chick, used in a simplified dilluent system for Japanese encephalitis virus haemagglutination—inhibition tests,

1,299

storage for use in haemagglutination assay, 1, 209

Fc fragment, of IgG, application of labelled serum for identification of anticytomegalovirus IgG and IgM antibodies, 1, 133

field-flow fractionation, of alkali-liberated nuclear polyhedrosis virus, 1, 241

field isolates, of rotavirus, analysis of genome profiles of, 2, 203

filter chromatography, concentration of bacteriophage lysates by, 1, 79

floating technique, method for virucide testing, 1, 27

Flury HEP, rabies strain, identified by use of monoclonal antibodies. 1.33

formalinised chick erythrocytes, used in the serological diagnosis of arboviruses, 1, 299

French beans, in immunochemical detection of ds RNA, 2, 277

fulminant viral hepatitis, 2, 17

gastroenteritis, rotavirus from infantile faeces, identified by electron microscopy, 2, 367

genome profiles, of field isolates of rotavirus, analysis of, 2, 203

genome, studied by the staining and shadowing method, 2, 159

gipsy moth (*Lymantria dispar* Linnaeus), fieldflow fractionation of alkali-liberated nuclear polyhedrosis virus from, 1, 241

glycoprotein, C-type viral, detection and quantification of by ELISA, 2, 357

growth inhibition, by interferon of cells in monolayer, measurement of, 1, 197

haemagglutination assay, storage of turkey erythrocytes for use in, 1, 209

haemagglutination—inhibition tests, for Japanese encephalitis virus, 1, 299

haemagglutination

immune adherence, 2, 31
passive, aldehyde-fixed erythrocytes coupled to HBsAg for use in, 2, 269

rosettes, in purified rubella virus, 2, 195 Hepanostika, for HBsAg, EIA for, 2, 57 hepatitis A epidemiology of in Amsterdam, ELISA for diagnosis of, 2, 47 past, present and future, 2, 1 pathogenesis and serodiagnosis of acute infection, 2, 31 hepatitis B clinical relevance of anti-HBc in, 2, 107 cross-reactivity between and non-A, non-B used to identify and detect non-A, non-B e antigen, 1, 149 HBeAg, anti-HBe and anti-HBc in patients with, 2, 97 Hepanostika, EIA for diagnosis of, 2, 57 past, present and future, 2, 1 post-transfusion, in the Blood Bank, 2, 119 virus, characterisation of, 1, 47 virus, marker of viral replication, 2, 7 virus-specific DNA polymerase, technique for the detection of, 2, 349 hepatitis B core antigen (HBcAg), 2, 7 comparison of RIA and counterimmunoelectrophoresis for the detection of, 2, 181 detection and significance of in the Blood Bank, 2, 119 hepatitis B e antigen (HBeAg) comparison of enzyme-labelled F(ab'), and IgG conjugates in an EIA for, 2, 85 in patients with hepatitis B, 2, 97 see also HBcAG, 2, 7 hepatitis non-A, non-B virus clinical aspects, 2, 17 HBeAg identified by use of the cross-reactivity between hepatitis B and, 1, 149 long incubation, identification and detection of, 2, 127 hepatitis B surface antibody in patients with hepatitis B, 2, 97 production of monoclonal antibodies against, by somatic cell hybrids, 1, 257 see also HBcAg and HBeAg, 2, 7 hepatitis B surface antigen (HBsAg) aldehyde-fixed erythrocytes coupled to for use in passive haemagglutinin, 2, 269 confirmation of specificity by neutralisation in immunoradiometric assay for, 1, 113 EIA for rapid detection of, 2, 63 Hepanostika, an EIA for, 2, 57

in the Blood Bank, 2, 119

cases of, 2, 17 status and significance of testing for, by EIA, 2, 71 virus-specific DNA polymerase, modified technique for detection of, 2, 349 hepatitis B surface antigen carriers anti-HBc detected in the sera of, 2, 107 detection of hepatitis B virus-specific DNA polymerase in, 2, 349 hepatitis B virions purified from the sera of, 1.47 persistence of anti-HBc IgM in, 2, 97 herpes simplex virus in study of single radial complement fixation, 2, 261 production of hybrid cell lines secreting antibodies to, 2, 223 virions and nucleocapsids, separation of on Percoll gradients, 1, 303 hybrid cell lines, secreting antibodies to herpes simplex, production of, 2, 223 hybridoma cells detection of monoclonal influenza antibodies by, 1, 275 secreting anti-HBs, 1, 257 IgG. anticy to megalovirus, assayed by radioimmunological method, 1, 133 anticytomegalovirus, demonstrated by an interchangeable ELISA, 2, 301, 315 anti-rubella, ELISA of, 1,1 conjugates, EIA for HBeAg, 2, 85 IgM anti-HBc, in patients with hepatitis B, 2, 97 anticytomegalovirus, assayed by radioimmunological method, 1, 133 anticytomegalovirus, demonstrated by an interchangeable ELISA, 2, 301 anti-rubella, ELISA of, 1, 1 rubella-specific, tested for by solid-phase immunosorbent haemadsorption test, 2. 341 immune adherence haemagglutination, for assay of larger numbers of specimens for HAV or anti-HAV, 2, 31

immunoradiometric assay for detection of,

-negative chronic active hepatitis, non-A,

non-B, an aetiological factor in some

immune electron microscopy

the use of markers in, 2, 169

solid phase, specific migration on the grid associated with, 1, 325

methods for the detection of HAV, 2, 31

immunoassays, for the detection of HAV, 2, 31

immunoperoxidase, anti-IgG, technique for detection of Epstein-Barr virus, 1, 11

immunoradiometric assay, see radioimmunoassay

immunosorbent haemadsorption, solid-phase, test for rubella-specific IgM, 2, 341

impression smears, of brains from infected animals, presence of antigens of rabies origin in, 1, 33

inclusion bodies, isolation and purification of, 1,241

indirect immune electron microscopy, for rapid identification of viruses, 2, 367

indirect immunofluorometric assay, solid-phase, detection of monoclonal influenza antibodies with, *I*, 275

infectious bronchitis virus, disruption of with Triton X-100 detergent, 1, 121

influenza virus

detection of previous antigenic priming with, 1, 355

in study of single radial complement fixation test, 2, 261

method for purification of human erythrocyte receptors for, 1, 285

monoclonal antibodies to, synthesised by hybridoma cells, 1, 275

plaque assay for in Madin-Darby canine kidney cells, 1, 117

separation of M-protein from by agarose gel electrophoresis, 2, 211

intact particles, of bacteriophage $\phi X174$ electron microscopic studies of, 2, 159

interferon

for antiviral therapy, 2, 7

method for measuring growth inhibition by, 1, 197

murine leukemia virus yield-reduction semimicroassay for, 1, 167

IUdR inhibition, cells made impermissive by, 2,321

Japanese encephalitis virus, haemagglutination inhibition tests, diluent system for, 1, 299

lectin, *Crotalaria juncea*, affinity chromatography of bovine viral diarrhea virus on, 2, 293

Lymantria dispar Linnaeus, gipsy moth, fieldflow fractionation of alkali-liberated nuclear polyhedrosis virus from, I, 241

lymphoblastoid B.95-8 cells, continuous culturing of in the presence of phosphonoacetic acid, 1, 349

lysates, bacteriophage, concentration of by filter chromatography, 1, 79

Madin-Darby canine kidney cells, simplified plaque assay for influenza viruses in, I, 11

mammalian RNA tumour viruses, production of in serum-free culture medium, allowing rapid RNA subunit purification, 1, 213

markers, use of immune electron microscopy, 2, 169

measles virus RNA, high-specificity radiolabelled 50S, preparation and characterisation of, 1, 223

metrizamide, in purification of infectious rubella virus, 2, 141

microassay, for the titration of human respiratory syncytial virus, 1, 343

monoclonal antibodies

against HBsAg, production of by somatic cell hybrids, 1, 257

CVS, rabies strains, identified by use of, 1, 33

to herpes simplex virus type 2, secreted by hybrid cell lines, 2, 223

to influenza, detection of with a solid-phase indirect immunofluorometric assay, 1, 275

Mount Elgon bat virus, direct plaque assay for, 1,201

M-protein, separation of from influenza virus by agarose gel electrophoresis, 2, 211

murine leukaemia virus yield-reduction semimicroassay for interferon, 1, 167

neonates with congenital cytomegalovirus, routine screening of, 2, 315

non-ionic detergent β-D-octylglucoside, in sucrose density gradient for characterisation of rubella virus, 2, 195

nuclear antigen, Epstein-Barr, characterisation of, 1, 11

nuclear polyhedrosis virus from gipsy moth, field-flow fractionation of, 1, 241

nuclear T-antigen, contained in transformed cells of SV40, 2, 331

nucleocapsids of herpes simplex virus, separated on Percoll gradients, 1, 303

nucleoprotein complexes, prepared from vaccinia virions irradiated with UV light, l, 185

 β-D-Octylglucoside, non-ionic detergent, for characterisation of rubella virus, 2, 195
 orthopoxvirus DNA, preparation of, 2, 175

papova virus, use in construction of somatic cell hybrids, 1, 139

parainfluenza, type 3-infected target cells, measurement of cytotoxic responses against, 2, 237

pathogenesis, of acute hepatitis A virus, 2, 31
Percoll gradients, separation of herpes simplex
virus virions and nucleocapsids on, 1,
303

phosphonoacetic acid, continuous culturing of B.95-9 cells in the presence of, 1, 349

plant viruses, comparison of two, 1, 179

plaque inhibition technique, for assay of antibodies to influenza virus, used to detect previous antigenic priming with influenza virus, 1, 355

polysome precipitation, antibody-mediated, as a method for size determination of viral mRNA species, 1, 99

pseudoreplica method, of electron microscopy, evaluated as a rapid screening technique for detection of cytomegalovirus in neonates, 2, 315

rabbit antisera, homologous and heterologous, for neutralisation of human adenovirus by CPE inhibition, 2, 321

rabies virus infection, diagnosis of by use of monoclonal antibodies, 1, 33

radioimmunoassay (RIA)

compared to counterimmunoelectrophoresis

for detection of antibody to HBcAg, 2, 181

compared with EIA for detection of human • rotaviruses and adenoviruses, 1, 331

for detection and differentiation of cytomegalovirus antibodies, 1, 133

for detection of HBsAg, 1, 311

for HBsAg, confirmation of specificity by neutralisation in, 1, 113

rheumatoid factor, interference of in ELISA of rubella IgM IgG antibodies, 1, 1

respiratory syncytial virus, human, microassay for the titration of, 1,343

ribavirin, antiviral activity of, 2, 7 RNA

ds, immunochemical detection of in plants, 2, 277

high-specific activity radiolabelled 50S measles, preparation and characterisation, 1, 223

messenger, size determination of by antibody-mediated polysome precipitation, 1, 99

³² P-labelling of viroid for hybridisation studies, 1, 229

subunit, rapid purification of, 1, 213 T₄ ligase, 2, 203

rosettes, rubella virus haemagglutinin, 2, 195 rotavirus

comparison of EIA and RIA for detection of from stool specimens, 1, 331

examined by immune electron microscopy for markers, 2, 169

field isolates of, method for analysing genome profiles of, 2, 203

from infantile gastroenteritis faeces, identified by immune electron microscopy, 2, 367

Rous sarcoma virus, purification of reverse transcriptase from, 1, 156

rubella

ELISA of, IgM and IgG antibodies, interference of IgM rheumatoid factor in, 1,

purification of infectious virus, comparison of two methods, 2, 141

-specific IgM, solid phase immunosorbent haemadsorption test for, 2, 341

virus haemagglutinin rosettes, characterisation of, 2, 195

scanning electron microscopy, for location and quantitation of antigen on the surface of virus-infected cells, 1, 61

screening of blood donors/recipients by improved EIA for HBeAg, 2, 85 for HBsAg and anti-HBs by EIA, 2, 71 in the Blood Bank, 2, 119

screening, routine, of newborns with congenital cytomegalovirus infection, by electron microscopy, 2, 315

semi-microassay, murine leukaemia virus yield-reduction, for interferon, 3, 167

Sepharose 2B

compared to gradient centrifugation for purification of infectious rubella virus, 2, 141

coupled to *Crotalaria juncea* for purification of bovine diarrhea virus by chromatography, 2, 293

for purification of reverse transcriptase, 1, 156

serodiagnosis, of acute hepatitis A virus, 2, 31 serum-free culture medium, production of mammalian RNA tumour viruses in, to allow rapid RNA subunit purification, 1, 213

silkworm *Bombyx mori*, use of electron microscopic studies of bacteriophage ϕ X174 intact and 'eclipsing' particles, 2, 159 simian virus 40,

replicating, super-coiled, detection of by electrophoresis, 1, 235

transformation of canine kidney cells by, 2,331

single radial complement fixation test, application to serological diagnosis of viruses, 2, 261

sodium metrizoate, in purification of infectious rubella virus, 2, 141

solid phase

competitive RIA for anti-HBc, 2, 181

immune electron microscopy, associated with SMOG, 1, 325

immunoradiometric assay for HBsAg, 1, 311

immunosorbent haemadsorption assay, for rubella-specific IgM, 2, 341

indirect immunofluorometric assay for detection of monoclonal influenza antibodies, 1, 275 RIA for determination of susceptibility of varicella, 2, 149

somatic cell hybrids

constructed with embryonic kidney cells, for study of viral gene expression, 1, 139

papova virus, use in construction of, 1, 139 production of monoclonal antibodies against HBsAg by, 1, 257

stool specimens

comparison of EIA and RIA for detection of human rotaviruses and adenoviruses from, 1, 331

in acute phase of hepatitis A, 2, 31

subviral particles, preparation of from vaccinia virions irradiated with UV, 1, 185

syncytium formation, titration of human respiratory syncytial virus by, 1, 343

tobacco mosaic virus

in detection of ds RNA in plants, 2, 277

in method of raising antibodies against plant viruses, 1, 179

tobacco ringspot virus, in detection of ds RNA in plants, 2, 277

T₄ RNA ligase, used to visualise genome RNA profiles of field isolates of rotavirus, 2, 203

transplant recipients, in non-A, non-B hepatitis patients, 2, 17

Triton X-100 detergent

for disruption of infectious bronchitis virus, 1, 121

treatment of purified rubella virus with, 2,

tumour viruses, mammalian RNA, production of allowing rapid RNA subunit purification, 1, 213

turkey erythrocytes, storage of for use in virus haemagglutination assays, 1, 209

ultrastructure and ultrastructural changes, non-A, non-B hepatitis infection, 2, 17

³ H-uridine, used in terminal labelling assay to measure cell-mediated cytotoxic responses against parainfluenza type 3-infected target cells, 2, 237

vaccinia virions, preparation of subviral particles from, 1, 185

varicella

determination of susceptibility to by solidphase RIA, 2, 149

lack of cross-reactivity with cytomegalovirus, 2, 301

Vero cells

measles virus-infected, used in hybridisation studies, 1, 223

neutralisation of adenovirus infectivity and cytotoxin studies, 1, 349

use in direct plaque assay for Mount Elgon bat virus, 1, 201

viral capsid antigen, Epstein-Barr virus, presence of tested for, 1, 349

viral cores, extracted from virus-infected cell cultures, for preparation of orthopoxvirus DNA, 2, 175

viral gene expression, studied in selection of

somatic cell hybrids, 1, 139

viral glycoprotein

contained in rubella virus haemagglutinin rosettes, 2, 195

C-type, detected and quantified by ELISA, 2, 357

virions and nucleocapsids of herpes simplex separated on Percoll gradients, 1, 303

viroid RNA, ³² P-labelling of, for hybridisation studies, *I*, 229

virucide testing, new method for, 1, 27

virus-specific DNA polymerase, to hepatitis B, detection of, 2, 349

WHO anti-HBs standard, for sensitivity comparisons in evaluation of RIA and EIA, 2,63

AUTHOR INDEX – Volumes 1 and 2

Alain, R., see Berthiaume, L., 2, 267

Aldershvile, J. and Nielsen, J.O.

HBeAg, anti-HBe and anti-HBc IgM in patients with hepatitis B, 2, 97

Almeida, J.D., Skelly, J., Howard, C.R. and Zuckerman, A.J.

The use of markers in immune electron microscopy, 2, 169

Almeida, J.D., Stannard, L.M. and Shersby, A.S.M.

A new phenomenon (SMOG) associated with solid phase immune electron microscopy, 1, 325

Altenburg-Petrie, B. and Mayor, H.D.

Growth of adeno-associated virus in embryonated eggs, 2, 287

Arnold, W., see Hess, G., 2, 107

Ashton, B.N., see Spruance, S.L., 1, 223

Babin, S., see Trepo, C., 2, 127

Bacchetti, S., see Killington, R.A., 2, 223

Bailey, P.D., see Macris, M.P., 2, 315

Balachandran, N., see Killington, R.A., 2, 223

Banatvala, J.E., see Goldwater, P.N., 2, 341

Barbara, J.S., see Cameron, C.H., 1, 311

Bar-Joseph, M. and Malkinson, M.

Hen egg yolk as a source of antiviral antibodies in the enzyme-linked immunosorbent assay (ELISA): a comparison of two plant viruses, *I*, 179

Bedarida, G., see Zanetti, A.R., 2, 71

Berthiaume, L., Micusan, V., Alain, R. and Trépanier, P.

Rapid identification of viruses by a simple indirect immune electron microscopy technique using ferritin-labelled antibodies, 2, 367

Bianchi, V., see Zanetti, A.R., 2, 71

Birbeck, T.H., see Liddell, K.G., 1, 209

Bos, E., see Van der Waart, M., 2, 85 Bossman, H.B., see Taylor, D.H., 2, 251

Bradley, D.W.

Hepatitis A virus infection: pathogenesis

and serodiagnosis of acute disease, 2, 31 Brandt, K.H., see Katchaki, J.N., 2, 119 Brann, A.W., see Macris, M.P., 2, 315 Briggs, M., see Cameron, C.H., 1, 113 Brouwer, R., see Katchaki, J.N., 2, 119 Bruijnis, F., see Van der Waart, N., 2, 85 Burness, A.T.H., see Pardoe, I.U., 1, 285

Caldwell, K.D., Nguyen, T.T., Giddings, J.C. and Mazzone, H.M.

Field-flow fractionation of alkali-liberated nuclear polyhedrosis virus from gypsy moth *Lymantria dispar* Linnaeus, 1, 241

Cameron, C.H. and Briggs, M.

Confirmation of specificity by neutralisation in immunoradiometric assay for hepatitis B surface antigen, 1, 113

Cameron, C.H., Combridge, B.S., Howell, D.R. and Barbara, J.A.J.

A sensitive immunoradiometric assay for the detection of hepatitis B surface antigen, 1, 311

Campbell-Benzie, A., Kangro, H.O. and Heath, R.B.

The development and evaluation of a solidphase radioimmunoassay (RIA) procedure for the determination of susceptibility to varicella, 2, 149

Chevallier, P., see Trepo, C., 2, 127

Clarke, I.N. and McCrae, M.A.

A rapid and sensitive method for analysing the genome profiles of field isolates of rotavirus, 2, 203

Cohen, B.J., Hewish, R.A. and Mortimer, P.P. Comparison of radioimmunoassay and counterimmunoelectrophoresis for the detection of antibody to hepatitis B core antigen, 2, 181

Combridge, B.S., see Cameron, C.H., 1, 311

Condit, R., see Esposito, J., 2, 175

Cote, P.J. Jr., see Shih, J.W.-K., 1, 257

Coutinho, R.A., Duermeyer, W. and Van der Veen, J.

Epidemiology of hepatitis A in Amsterdam, October 1978 – December 1979, 2, 47 Croese, J., see Wolters, G., 2, 57

Croteau, G., see Lecomte, J., 2, 211

D'Agostino, F., see Zanetti, A.R., 2, 71
Dapolito, G.M., see Shih, J.W.-K., 1, 257
Dodd, R.Y., see Fang, C.T., 2, 349
Duman, R.G., see Gaush, C.R., 2, 331
Duermeyer, W., see Coutinho, R.A., 2, 47

Ehlke, S., see Major, E.O., I, 139
El Mekki, A.A., see Van der Groen, G., I, 27
Erfle, V., see Schetters, H., 2, 357
Esposito, J., Condit, R. and Obijeski, J.
The preparation of orthopoxvirus DNA, 2, 175

Fang, C.T., Nath, N., Pielech, M. and Dodd, R.Y.

A modified technique for the detection of hepatitis B virus-specific DNA polymerase, 2,349

Feorino, P.M., see Phillips, D.J., 1, 275
Ferroni, P., see Zanetti, A.R., 2, 71
Finke, J.H., see Hsu, L., 1, 167
Flamand, A., see Wiktor, T.J., 1, 33
Francki, R.I.B., see Gould, A.R., 2, 277
Friedrich, T.D. and Lehman, J.M.

An electrophoretic method for the detection of replicating, supercoiled simian virus 40 DNA, 1, 235

Gaush, C.R., Duman, R.G. and Hougland, A.E. Transformation of canine kidney cells by SV40, 2, 331

Gerba, C.P., see Goyal, S.M., 1, 79

Gerin, J.L., see Shih, J.W.-K., 1, 47, 257

Gibbins, B.L., see Williamson, S.I., 2, 237 Giddings, J.C., see Caldwell, K.D., 1, 241

Goldblum, N., see Margalith, M., 1, 349

Goldwater, P.M. and Banatvala, J.E.

Solid-phase immunosorbent haemadsorption (SPIHAd): a new, rapid and simple test for rubella-specific IgM, 1, 341

Golomb, M., Vora, A.C. and Grandgenett, D.P.
 Purification of reverse transcriptase from avian retroviruses using affinity chromatography on heparin-Sepharose, 1, 157
 Gomwalk, N.E.

A direct plaque assay for Mount Elgon bat virus, 1, 201

Gould, A.R. and Francki, R.I.B.

Immunochemical detection of ds-RNA in healthy and virus-infected plants and specific detection of viral ds-RNA by hybridization to labelled complementary DNA, 2, 277

Goyal, S.M., Zerda, K.S. and Gerba, C.P.
Concentration of bacteriophage lysates by filter chromatography, 1, 79
Grandgenett, D.P., see Golomb, M., 1, 157

Grandgenett, D.P., see Golomb, M., 1, 157 Grimaud, J.A., see Trepo, C., 2, 127 Grill, L.K., see Negruk, V.I., 1, 229 Gut, W., see Jankowski, M.A., 1, 133

Haas, H. and Hotz, G.

Rapid detection of HBsAg and anti-HBs by enzyme-immunoassay, 2, 63

Halonen, P., see Sarkkinen, H.K., 1, 331

Hantz, O., see Trepo, C., 2, 127

Hantz, O., see Vitvitski, L., 1, 149

Hayashi, K., see Niwayama, S., 2, 261

Heath, R.B., see Campbell-Benzie, A., 2, 149

Hehlmann, R., see Schetters, H., 2, 357

Hess, G. see Shih, J.W.-K., 1, 47

Hess, G. and Arnold, W.

The clinical relevance of the antibody to hepatitis B core antigen (anti-HBc): a review, 2, 107

Hewish, R.A., see Cohen, B.J., 2, 181

Holowczak, J.A., see Soloski, M., 1, 185

Hotz, G., see Haas, H., 2, 63

Hougland, A.E., see Gaush, C.R., 2, 331

Howard, C.R., see Almeida, J.D., 2, 169

Howard, C.R., see Lancer, J.A., 1, 121

Howell, D.R., see Cameron, C.H., 1, 311

Hsu, L., Finke, J.H. and Proffitt, M.R.

A murine leukemia virus yield-reduction semi-microassay for interferon, 1, 167 Huprikar, J.A. and Rabinowitz, S.

A simplified plaque assay for influenza viruses in Madin-Darby kidney (MDCK) cells, 1, 117

Ikram, H. and Prince, A.M.

A method for coupling the hepatitis B surface antigen to aldehyde-fixed erythrocytes for use in passive hemagglutination, 2, 269 Ito, H. and Iwasa, S.

Simple serological diagnosis of arboviruses:

a simplified diluent system for Japanese encephalitis virus hemagglutination—inhibition tests using formalinized chick erythrocytes, 1, 299

Iwasa, S., see Ito, H., 1, 299 Iwase, I., see Niwayama, S., 2, 261

Jankowski, M.A., Gut, W., Nawrocka, E. and Kańtoch, M.

Detection and differentiation of cytomegalovirus antibodies by radioimmuno-assay, 1, 133

Jeansson, T., see Svennerholm, B., 1, 303

Kangro, H.O., see Campbell-Benzie, A., 2, 149 Kaplan, P.M., see Shih, J.W.-K, 1, 47 Karsnas, P., see Moreno-Lopez, J., 2, 293

Katchaki, J.N., Siem, T.H., Brouwer, R., Brandt, K.H. and Van der Waart, M.

Detection and significance of anti-HBc in the Blood Bank; preliminary results of a controlled prospective study, 2, 119

Katchaki, J., see Van der Waart, M., 2, 85 Kańtoch, M., see Jankowski, M.A., 1, 133 Katz, E., see Zakay-Rones, Z., 1, 355

Kendal, A.P., see Phillips, D.J., 1, 275

Killington, R.A., Newhook, L., Balachandran, N., Rawls, W.E. and Bacchetti, S.

Production of hybrid cell lines secreting antibodies to herpes simplex virus type 2, 2, 223

Kimball, P.C., see Strauss, E.M., 1, 213 Koprowski, H., see Wiktor, T.J., 1, 33 Kristiansen, T., see Moreno-Lopez, J., 2, 293

Kuipers, L.P.C., see Wolters, G., 2, 57 Kurstak, E., see Van den Hurk, J., 1, 11

Kurth, R., see Netuschil, L., 1, 99

Lampert, M., see Major, E.O., 1, 139 Lancer, J.A. and Howard, C.R.

The disruption of infectious bronchitis virus (IBV-41 strain) with Triton X-100 detergent, *I*, 121

Lecomte, J. and Croteau, G.

Separation of M-protein from influenza virus by agarose gel electrophoresis, 2, 211 Lee, F.K., see Macris, M.P., 2, 315

Lehman, J.M., see Friedreich, T.C. 1, 235 Levy, R., see Zakay-Rones, Z., 1, 355

Liddell, K.G. and Birkbeck, T.H.

Storage of turkey erythrocytes for use in virus haemagglutination assays, 1, 209 Logan, K.B., Rees, G.E., Seeley, N.D. and

Primrose, S.B.

Rapid concentration of bacteriophages from large volumes of freshwater: evaluation of positively charged, microporous filters, I, 87

Lundén, R., see Svennerholm, B., 1, 303 Lycke, E., see Svennerholm, B., 1, 303

Macris, M.P., Nahmias, A.J., Bailey, P.D., Lee, F.K., Visintine, A.M. and Brann, A.W.

Electron microscopy in the routine screening of newborns with congenital cytomegalovirus infection. 2, 315

Major, E.O., Ehlke, S. and Lampert, M.
Selection of somatic cell hybrids between
BK virus transformed BHK-21 and human
embryonic kidney cells to study viral gene
expression, 1, 139

Malkinson, M., see Bar-Joseph, M., 1, 179 Manor, D., see Margalith, M., 1, 349

Marchessault, F., see Trudel, M., 2, 141, 195
Margalith, M., Manor, D. and Goldblum, N.
On the continuous culturing of B.95-8
cells in the presence of phosphonoacetic
acid, 1, 349

Margalith, M., see Zakay-Rones, Z., 1, 355 Mayor, H.D., see Altenburg-Petrie, B., 2, 287 Mazzone, H.M., see Caldwell, K.D., 1, 241 McCrae, M.A., see Clarke, I.N., 2, 203

Micusan, V., see Berthiaume, L., 2, 367

Moreno-Lopez, J., Kristiansen, T. and Karsnas, P.

Bovine viral diarrhea virus: affinity chromatography on *Crotalaria juncea* lectin, 2, 293

Mortimer, P.P., see Cohen, B.J., 2, 181 Motoda, S., see Niwayama, S., 2, 261

Nahmias, A.J., see Macris, M.P., 2, 315 Nath, N., see Fang, C.T., 2, 349

Nawrocka, E., see Jankowski, M.A., 1, 133
 Negruk, V.I., Grill, L.K. and Semancik, J.S.
 In vitro ³²P-labelling of viroid RNA for hybridization studies, 1, 229

Netuschil, L. and Kurth, R.

Antibody-mediated polysome precipitation as a method for the size determination of

viral mRNA species: viral envelope glycoprotein mRNA of avian sarcoma viruses, 1,99

Newhook, L., see Killington, R.A., 2, 223 Nguyen, T.T., see Caldwell, K.D., 1, 241

Nielsen, J.O., see Aldershvile, J., 2, 97

Niwayama, S., Sato, S., Ochiai, H., Motoda, S., Iwase, I. and Hayashi, K.

Study on the single radial complement fixation (SRC-FIX) test, 2, 261

Obijeski, J., see Esposito, J., 2, 175 Ochiai, H., see Niwayama, S., 2, 261 Olofsson, S., see Svennerholm, B., 1, 303

Payment, P., see Trépanier, P., 1, 343

Pardoe, I.U. and Burness, A.T.H.
Preparation of erythrocyte receptors for viruses by affinity chromatography, 1, 285
Parry, J.E., see Pringle, C.R., 1, 27
Pattyn, S.R., see Van der Groen, G., 1, 27

Payment, P., see Trudel, M., 2, 141, 195 Phillips, D.J., Kendal, A.P., Webster, R.G., Feorino, P.M. and Reimer, C.B.

Detection of monoclonal influenza antibodies synthesized in culture by hybridoma cells with a solid-phase indirect immunofluorometric assay, 1, 275.

Pichoud, C., see Trepo, C., 2, 127

Pielech, N., see Fang, C.T., 2, 349

Primrose, S.B., see Logan, K.B., 1, 87

Prince, A.M., see Ikram, H., 2, 269

Pringle, C.R. and Parry, J.E.

Location and quantitation of antigen on the surface of virus-infected cells by specific bacterial adherence and scanning electron microscopy, 1, 61

Prins-Bekius, M., see Van der Waart, M., 2, 85

Profitt, M.R., see Hsu, L., 1, 167

Rabinowitz, A., see Huprikar, J., I, 117 Rawls, W.E., see Killington, R.A., 2, 223 Rees, G.E., see Logan, K.B., I, 87 Reimer, C.B., see Phillips, D.J., I, 275

Sarkkinen, H.K., Tuokko, H. and Halonen, P.E. Comparison of enzyme-immunoassay and radioimmunoassay for detection of human rotaviruses and adenoviruses from stool specimens, 1, 331

Sato, S., see Niwayama, S., 2, 261

Schellekens, H. and Stitz, L.W.

Simple method for measuring growth inhibition by interferon of cells in monolayer, 1, 197

Schetters, K., Hehlmann, R. and Erfle, V.
ELISA for the detection and quantification of C-type viral glycoprotein (gp70) using antibodies that recognize the protein moieties of the glycoproteins, 2, 357

Schrader, E. and Wigand, R.

Neutralization of adenovirus infectivity and cytotoxin in various cell cultures, 2, 321

Schuurs, A., see Van der Waart, M., 2, 85

Seeley, N.D., see Logan, K.B., 1, 87

Semancik, J.S., see Negruk, V.I., 1, 229

Septjan, M., see Trepo, C., 2, 127

Shersby, A.S.M., see Almeida, J.D., 1, 325 Shih, J.W.-K., Cote, P.J. Jr., Dapolito, G.M. and Gerin, J.L.

Production of monoclonal antibodies against hepatitis B surface antigen (HBsAg) by somatic cell hybrids, 1, 257

Shih, J.W.-K., Hess, G., Kaplan, P.M. and Gerin, J.L.

Characterization of hepatitis B virus (Dane particle), 1,47

Siem, T.H., see Katchaki, J.N., 2, 119

Simon, M.C., see Strauss, E.M., I, 213

Skelly, J., see Almeida, J.D., 2, 169

Smith, C.B., see Spruance, S.L., 1, 223

Snelting, A., see Van der Waart, M., 2, 85

Soloski, M. and Holowczak, J.A.

Preparation of subviral particles from vaccinia virions irradiated with ultraviolet light, 1, 185

Spruance, S.L., Ashton, B.N. and Smith, C.B.
Preparation and characterization of highspecific activity radiolabeled 50S measles
virus RNA, 1, 223

Stannard, L.M., see Almeida, J.D., 1, 325

Stitz, L.W., see Schellekens, H., 1, 197

Strauss, E.M., Simon, M.C. and Kimball, P.C. Efficient production of mammalian RNA tumor viruses in scrum-free culture medium allows rapid RNA subunit purification, 1, 213

Sundqvist, V.-A. and Wahren, B.

An interchangeable ELISA for cytomegalo-

virus antigen and antibody, 2, 301

Svantesson, G., see Svennerholm, B., 1, 303 Svennerholm, B., Vahlne, A., Jeansson, S.,

Lundén, R. Olofsson, S., Svantesson, G. and Lycke, E.

Separation of herpes simplex virus virions and nucleocapsids on Percoll gradients, I, 303

Taylor, D.H. and Bosmann, H.B.

Measurement of electrokinetic properties of vaccinia and reovirus by laser-illuminated whole-particle microelectrophoresis, 2, 251

Trépanier, P., Payment, P. and Trudel, M.

A simple and rapid microassay for the titration of human respiratory syncytial virus, 1, 343

Trépanier, P., see Berthiaume, L., 2, 367

Trepo, C., Vitvitski, L., Hantz, O., Chevallier, P., Pichoud, C., Babin, S., Grimaud, J.A. and Sepetjan, M.

Identification and detection of long-incubation non-A, non-B hepatitis virus and associated antigens or antibodies, 2, 127

Trepo, C., see Vitvitski, L., 1, 149

Trudel, M., see Trépanier, P., 1, 343

Trudel, M., Marchessault, F. and Payment, P. Purification of infectious rubella virus by gel filtration on Sepharose 2B compared to gradient centrifugation in sucrose, sodium metrizoate and metrizamide, 2, 141

Trudel, M., Marchessault, F. and Payment, P. Characterisation of rubella virus hemagglutinin rosettes, 2, 195

Tuokko, H., see Sarkkinen, H.K., 1, 331

Vahlne, A., see Svennerholm, B., 1, 303 Van den Hurk, J. and Kurstak, E.

Characterization of Epstein-Barr nuclear antigen (EBNA). I. A new technique for the detection of EBNA or anti-EBNA anti-bodies and its applicability to the study of chromosome-EBNA interactions, *I*, 11

Van der Groen, G., El Mekki, A.A. and Pattyn, S.R.

New method for virucide testing, 1, 27 Van der Veen J., see Coutinho, R.A., 2, 47 Van der Waart, M., see Katchaki, J.N., 2, 119 Van der Waart, M., Snelting, A., Bruijnis, F., Prins-Bekius, M., Bos, E., Katchaki, J. and Schuurs, S.

Comparison of enzyme-labelled $F(ab')_2$ and IgG conjugates in an enzyme-immuno-assay for hepatitis B 'e' antigen, 2, 85

Vejtorp, M.

The interference of IgM rheumatoid factor in enzyme-linked immunosorbent assays of rubella IgM and IgG antibodies, 1, 1

Visintine, A.M., see Macris, M.P., 2, 315

Vitvitski, L., Trepo, C. and Hantz, O.

Use of the cross-reactivity between hepatitis B and non-A, non-B viruses for the identification and detection of non-A, non-B 'e' antigen, 1, 149

Vitvitski, L., see Trepo, C., 2, 127

Vogten, A.M.

Viral hepatitis B: the past, the present and the future, 2, 1

Vora, A.C. see Golomb, M., 1, 157

Wahren, B., see, Sundqvist, V.-A., 2, 301

Webster, R.G., see Phillips, D.J., 1, 275

Wigand, R., see Schrader, E., 2, 321

Wiktor, T.J., Flamand, A., and Koprowski, H.
Use of monoclonal antibodies in diagnosis
of rabies virus infection and differentiation
of rabies and rabies-related viruses, 1, 33

Williams, R., see Wyke, R.J., 2, 17 Williamson, S.I. and Gibbins, B.L.

A terminal labelling assay using ³ H-uridine to measure cell-mediated cytotoxic responses against parainfluenza type 3-infected target cells, 2, 237

Wolters, G., Kuijpers, L.P.C. and Croese, J. Improved Hepanostika, an enzyme-immuno-assay for hepatitis B surface antigen, 2, 57 Wyke, R.J. and Williams, R.

Clinical aspects of non-A, non-B hepatitis infection, 2, 17

Yazaki, K.

Electron microscopic studies of bacteriophage ϕ X174 intact and 'eclipsing' particles, and the genome by the staining and shadowing method, 2, 159

Zakay-Rones, Z., Margalith, M., Levy, R. and Katz, E.

A sensitive plaque inhibition technique for assay of antibodies to influenza virus: use

to detect previous antigenic priming with influenza viruses, 1, 355

Zanetti, A.R., Bedarida, G., Ferroni, P., D'Agostino, F. and Bianchi, V.
Status and significance of testing for hepatitis B surface antigen and surface antibody, 2, 71 Zerda, C.S., see Goyal, S.M., 1, 79 Zuckerman, A.J.

Introduction to the virology of human hepatitis, 2, 7

Zuckerman, A.J., see Almeida, J.D., 2, 169



